

## Spectral Measures

Victor Guillemin, MIT

### Abstract

**Lecture 1.** Let  $X$  be a compact Riemannian manifold and let  $f_n$ ,  $n = 1, 2, \dots$ , be the eigenfunctions of the Laplace operator, normalized so that  $\int |f_n|^2 dx = 1$ . If  $X$  is negatively curved, then, by a theorem of Shnirelman–Zelditch–Colin de Verdiere, the measures,

$$\mu_i = |f_{n_i}|^2 dx$$

tend weakly to  $dx$  as  $i$  tends to infinity for “most” sequences of integers,  $n_1, n_2, \dots$ . For Riemannian manifolds in general no such result seems to be true, however in some instances, because of the presence of symmetries or for other geometric reasons, the eigenvalues of  $\Delta$  tend to group into clusters

$$\lambda_{i,n}, \quad 1 \leq i \leq N(n)$$

and the averages of the corresponding measures

$$(*) \quad \frac{1}{N(n)} \sum \mu_{n_i}$$

have, in some of these cases, interesting asymptotic properties. One example of this is when  $X$  is a *Zoll* manifold. A *Zoll* manifold has, by definition, the property that the eigenvalues of  $\sqrt{\Delta}$  lie on bands

$$an + b - O\left(\frac{1}{n}\right) < \sqrt{\lambda_{i,n}} < an + b + O\left(\frac{1}{n}\right)$$

and in this lecture I will describe some beautiful results of Colin de Verdiere from the mid-eighties on the asymptotics of the measures  $(*)$  associated with these bands.

**Lecture 2.** One of Colin’s results is that for *Zoll* manifolds the function,  $N(n)$ , is a polynomial. Boutet de Monvel and I were able to obtain an explicit formula for this polynomial by noting some similarities between *Zoll* manifolds and complex projective varieties. We also discovered that Colin’s results have analogues for projective varieties: If  $X$  is a non-singular projective variety and  $\mathbb{L} \rightarrow X$  its canonical line bundle one can associate a spectral measure to  $\mathbb{L}^n$  by setting

$$\mu_n(f) = \text{trace } \pi_n M_f$$

where, for  $f \in C^0(X)$ ,

$$M_f : \Gamma_{\text{hol}}(\mathbb{L}^n) \rightarrow L^2(\mathbb{L}^n)$$

is multiplication by  $f$  and

$$\pi_n : L^2(\mathbb{L}^n) \rightarrow \Gamma_{\text{hol}}(\mathbb{L}^n)$$

is orthogonal projection and one gets an asymptotic expansion of  $\mu_n$  in inverse powers of  $n$  as  $n$  tends to infinity. In general this asymptotics isn’t as explicitly computable as in Colin’s case except in a few rare instances. One such instance is if  $X$  is a toric variety and  $\Delta$  its defining polytope. I will show that in this case the asymptotics of  $\mu_n$  is described by an Euler–Maclaurin formula for  $\Delta$ .